

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : G01N 33/53, C07K 16/00	A1	(11) International Publication Number: WO 00/14536 (43) International Publication Date: 16 March 2000 (16.03.00)
(21) International Application Number: PCT/US99/19597 (22) International Filing Date: 26 August 1999 (26.08.99) (30) Priority Data: 09/148,712 4 September 1998 (04.09.98) US (71) Applicant: NEW ENGLAND BIOLABS, INC. [US/US]; 32 Tozer Road, Beverly, MA 01915 (US). (72) Inventors: TAN, Yi; 2 Heath Circle, Lynnfield, MA 01940 (US). COMB, Michael, J.; 10 Eaglehead Road, Manchester, MA 01944 (US). (74) Agent: WILLIAMS, Gregory, D.; New England Biolabs, Inc., 32 Tozer Road, Beverly, MA 01915 (US).		(81) Designated States: CA, JP, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: PRODUCTION OF MOTIF-SPECIFIC AND CONTEXT-INDEPENDENT ANTIBODIES USING PEPTIDE LIBRARIES AS ANTIGENS (57) Abstract A method is provided for producing motif-specific, context-independent antibodies which are specific to at least one fixed amino acid residue, whether modified or unmodified, in the context of variable surrounding amino acid or peptide sequences. The method includes the steps of (1) constructing a peptide library featuring at least one fixed amino acid and variable surrounding amino acids, and (2) immunizing a host with this peptide library. Antibodies may optionally be isolated and purified from the antisera of this immunized host. The disclosed method encompasses both modified motifs, such as phosphothreonine, phosphoserine, MAPK consensus recognition sites, 14-3-3 consensus recognition sites, CDK consensus recognition sites, and acetylated-lysine, as well as unmodified motifs.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BR	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

**PRODUCTION OF MOTIF-SPECIFIC AND CONTEXT-
INDEPENDENT ANTIBODIES USING PEPTIDE LIBRARIES
AS ANTIGENS**

5 BACKGROUND OF THE INVENTION

The present invention relates to the production of motif-specific, context-independent antibodies which are specific to at least one fixed amino acid residue in the
10 context of variable surrounding amino acid or peptide sequences. Antibodies with these properties are useful in characterizing various forms of cellular regulation as well as serving to profile genome wide changes in cellular protein levels and protein modification.

15 Identifying the targets of intracellular signaling cascades are of major importance in understanding cell growth, differentiation, and cell death. Protein kinase cascades relay information from the cell surface to multiple
20 cellular compartments including the nucleus and more distant cell processes such as synapses (Karin et al., Curr. Opin. Cell. Biol. 6:415-424 (1994)). Although a few targets of protein phosphorylation have been identified, most remain unknown, particularly those that regulate cell growth and
25 differentiation. For example, the MAP kinase cascade is known to play an important role in the regulation of cell growth (Lewis et al., *Adv. Cancer Res.* 74:49-139 (1998), Crowley et

al., *Cell* 77:841-852 (1994)). However, beyond a handful of substrates, few protein targets responsible for the diverse actions of the MAP kinase cascade have been identified (Fukunaga and Hunter, *EMBO* 16(8):1921-1933 (1997),
5 Stukenberg et al., *Curr. Biol.* 7:338-348 (1997)).

Another example of cell signaling proteins are the 14-3-3 proteins, which represent a phylogenetically conserved family of phosphoserine binding proteins whose precise role in
10 cell signaling has yet to be determined (Burbelo and Hall, *Curr. Biol.* 5(2):95-96 (1995)). These proteins represent a large fraction of total brain protein and are known to bind a wide variety of signaling molecules including: ras, raf, bad, cdc25, and many others (Yaffe et al., *Cell* 91:961-971 (1997)).
15 Recently, it has been shown that 14-3-3 proteins bind specifically to phosphorylated sites on proteins with the following motif: RXRSXS*XP where S* is phosphoserine and X represents any amino acid (Muslin et al., *Cell* 84:889-897 (1996), Yaffe et al. *supra*(1997)).

20

Similarly, histones have long been known to be modified by acetylation at specific lysine residues. Acetylation of lysine in histones is thought to reduce protein-DNA interactions and serve to open chromatin in regions undergoing
25 transcription (Struhl, *Genes & Development*, 12:599-606 (1998)). Recently, other proteins associated with transcription complexes have been shown to be acetylated on

lysine although the functional significance is unclear (Imhof et al., *Curr. Biol.* 7:689-692 (1997), Struhl *supra* (1998)).

Antibodies against phosphotyrosine have proven to be of
5 great value in identifying and characterizing intracellular
signaling mechanisms (Ross et al., *Nature* 294:654 (1981),
Kozma et al., *Method. Enzymol.* 201:28 (1991), White and
Backer, *Method. Enzymol.* 201:65 (1991), Kamps, *Method.*
Enzymol. 201:101 (1991)). Their value derives from two
10 properties; 1) their ability to discriminate whether or not a
protein is tyrosine phosphorylated, and 2) their ability to
react with a large variety of different proteins. These
properties have proven invaluable in tracing intracellular
signaling pathways and identifying new targets of activated
15 tyrosine kinases.

Ideally, the most useful phosphotyrosine antibodies
should be as general as possible, that is they should recognize
phosphotyrosine independently of the protein sequences in
20 which it is embedded (context independent) so as to allow
detection of all possible phosphotyrosine residues. The most
successful approaches for producing phosphotyrosine
antibodies have utilized phosphotyrosine or phosphotyramine
coupled via their free amino groups to keyhole limpet
25 hemocyanin using hetero- or bifunctional crosslinking agents
(Frackelton et al., *Method. Enzymol.* 201:79 (1991), White and
Backer *supra* (1991), Wang, *Method. Enzymol.* 201:53 (1991),

- Kamps *supra* (1991)). Although currently produced polyclonal and monoclonal phosphotyrosine antibodies do recognize many different proteins, they often show crossreactivity with other phosphate containing compounds, for example,
- 5 mononucleotides (Frackelton et al. *supra* (1991), Kamps *supra* (1991)). More importantly, most phosphotyrosine antibodies raised in this fashion display variable sequence reactivity, depending not only on the phosphorylated amino acid, but also upon the amino acid sequences surrounding phosphotyrosine.
- 10 For example, the present inventors have observed that most phosphotyrosine antibodies do not recognize phosphotyrosine preceded by proline as found in the activation loop of JNK and hence do not react significantly with activated (tyrosine phosphorylated) JNK [(Tan et al. unpublished observations)].
- 15 The reason for the variable reactivity is likely due to the fact that the phosphotyrosine antigen is not presented directly to the immune system in the context of variable surrounding amino acids, but is instead presented as a hapten, inappropriately coupled to the KLH carrier via artificial
- 20 linkages. This approach tends to produce antibodies that react well with phosphotyrosine but are sometimes blocked by surrounding amino acids as they are not present in the antigen.

- Other approaches have utilized total cellular
- 25 phosphotyrosine containing proteins as immungens (Glenney, *Method. Enzymol.* 201:92 (1991), Wang *supra* (1991)) with considerable success but the context-dependence of the

resulting antibody specificities was not carefully determined, although antibodies raised in this fashion did react with a majority of tyrosine phosphorylated proteins. Estimates as to the fraction of tyrosine phosphorylated proteins detected
5 range from 50% to 94% (Kamps *supra* (1991)).

Attempts to use the above mentioned techniques to produce similar antibodies for phosphoserine and phosphothreonine have met with limited success. Antibodies
10 produced to date have limited crossreactivity and lower affinity for phosphoserine or phosphothreonine probably due to the poor immunogenicity of these phospho-amino acids compared to phosphotyrosine (Heffetz et al., *Method. Enzymol.* 201:44 (1991)). Context-dependence and low affinity have
15 limited the utility of currently available phosphoserine and phosphothreonine antibodies, especially when compared to phosphotyrosine antibodies

Site-specific phosphoserine and phosphothreonine
20 antibodies were first described by Nairn et al. in 1982 and have proven to be highly useful tools to study protein phosphorylation (Czernik et al., *Method. Enzymol.* 201:264 (1991), Czernik et al., *Neuroprot.* 6:56-61 (1995)). One drawback of this type of antibody is that a different antibody
25 needs to be produced for each site of interest. Clearly, development of antibodies that detect phosphoserine or phosphothreonine in a context-independent fashion would be

desirable for use in tracing serine/threonine kinase cascades and in defining their biological responses. Likewise, development of context-independent phosphotyrosine antibodies would overcome the limitations of currently
5 available antibodies.

Motif-specific, context-independent antibodies would also be useful in identifying new targets of 14-3-3 action (i.e., other proteins phosphorylated at this motif) and in
10 characterizing the protein kinases that phosphorylate these sites. Likewise antibodies reactive against acetylated lysine would serve as useful tools to study the functional significance of acetylation of histones.

15 Such antibodies can further be used as general reagents for detecting phosphorylation or other enzymatic modification *in vitro*, such as in high throughput kinase assays for drug screens, as a single antibody can be used to recognize many different phosphorylated substrates. Phosphotyrosine
20 antibodies are currently employed in high throughput kinase assays to screen for selective, high affinity tyrosine kinase inhibitors. Compounds or drugs that block enzyme activity are detected by their ability to inhibit kinase activity as determined by a reduction of phosphotyrosine antibody binding
25 to phosphorylated substrate. Similar assays can be set up to screen for pharmaceutically useful compounds using antibodies produced as described above for phosphoserine,

phosphothreonine, or antibodies detecting other protein modifications.

Antibodies that detect short motifs in a context-independent fashion will also be particularly useful in profiling genome wide changes in protein levels and protein modification. For example, the use of context-independent phosphothreonine antibodies and 2D gel electrophoresis to profile genome wide changes in protein phosphorylation (Patterson and Garrels, Cell Biology: A Laboratory Handbook 249-257 (1994), Academic Press) as the result of drug treatment or overexpression of a particular protein will undoubtedly prove useful in identifying potential drug-protein interactions and suggest new downstream targets for overexpressed proteins.

SUMMARY OF THE INVENTION

In accordance with the present invention, there is provided a method of producing antibodies that selectively recognize specified short amino acid motifs independent of the surrounding amino acid, peptide, or protein sequences. The method allows the production of antibodies that recognize modified single amino acids, for example phosphorylated serine, threonine, and tyrosine, or acetylated lysine, as well other unmodified or modified motifs of one or more amino acids.

The method encompasses the production and purification of highly context-independent antibodies that recognize specific and highly degenerate amino acid motifs, such as
5 those found in kinase consensus sequences or other enzyme binding sites. Furthermore, the method can be used to produce highly context-independent polyclonal or monoclonal antibodies.

10 Antibodies produced by the method of the present invention may be specific to virtually any protein motif, either modified or unmodified. For example, the method can be used to produce antibodies recognizing phosphothreonine alone or phosphothreonine in the context of several fixed amino
15 acids as found in the MAPK, 14-3-3, or cdk consensus sites. It can also be used to produce antibodies specific for other modified amino acids, for example, acetylated lysine, or to detect any short motif of one or more amino acids, in a context-independent fashion.

20

The present invention further provides a method of profiling large and diverse protein populations on a genome-wide scale by utilizing motif-specific, context-independent antibodies against motifs conserved on such proteins. For
25 example, phosphorylation-specific antibodies allow genome-wide profiling of changes in protein phosphorylation as a result of drug treatment.

The present invention also provides a method of identifying an unknown substrate of a known enzyme through the use of motif-specific, context-independent antibodies which are raised against motifs common to other substrates of the enzyme.

The use of such motif-specific, context-independent antibodies as a reagent for the detection of enzymatic modifications of a given motif within a substrate is also encompassed by the present invention.

BRIEF DESCRIPTION OF DRAWINGS

Figure 1a is a table depicting the specificity of the affinity-purified, polyclonal antibodies produced against a phosphorylated threonine peptide library in Example I, when tested against specific peptides.

Figure 1b is a table depicting the specificity of the phosphothreonine antibodies of Example I when tested against various phosphopeptide libraries.

Figure 1c is a Western analysis which depicts the reactivity of the phosphothreonine antibodies of Example I against cell extracts from cells treated with and without okadaic acid and against other phosphoproteins.

Figure 1d is a table depicting the context-independence of the anti- phosphothreonine antibodies of Example I as shown by immobilized grid.

5

Figure 2a is a table depicting the specificity of the affinity-purified, polyclonal antibodies produced against a phosphorylated PXS*P peptide library in Example II.

10 Figure 2b is a Western analysis depicting the reactivity of the phospho-PXS*P antibodies of Example II against cell extracts from cells treated with and without okadaic acid and against other phosphoproteins.

15 Figure 3a is a table depicting the lack of reactivity of the affinity-purified, polyclonal 14-3-3 antibodies of Example III when tested against non-phosphopeptides or phosphopeptides lacking the motif.

20 Figure 3b is a Western analysis depicting the reactivity of the phospho-14-3-3 antibodies of Example III against cell extracts from cells transfected with GST-Bad and with TPA.

25 Figure 4a is a table depicting the specificity of the monoclonal antibodies produced against a phosphorylated PXT*PXR library in Example IV.

Figure 4b is a Western analysis depicting the reactivity of the CDK consensus site monoclonal antibodies of Example IV against phosphorylated and nonphosphorylated RB protein.

5 Figure 5a is a Western analysis depicting the specificity of the acetylated-lysine antibodies of Example V against acetylated BSA.

10 Figure 5b is a Western analysis depicting the reactivity of the acetylated-lysine antibodies of Example V against various proteins present in C6-cell extracts when antibodies are preincubated with nonacetylated peptide library.

15 Figure 5c is a Western analysis depicting the reactivity of the acetylated-lysine antibodies of Example V against various proteins present in C6-cell extracts when antibodies are preincubated with acetylated peptide library.

20 Figure 5d is a Western analysis depicting the reactivity of the acetylated-lysine antibodies of Example V against the control acetylated BSA when antibodies are preincubated with acetylated peptide library.

DETAILED DESCRIPTION

25

The present invention is based upon the concept that the concentration of any individual sequence in a peptide library

used as antigen is extremely low and hence will be insufficient to drive an immune response in a host. The only antigenic determinants of sufficiently high concentration to drive the immune response are thus the fixed residues common
5 to each sequence, as well as the peptide backbone itself.

Immunizing a host with peptide libraries representing all 20 amino acids at each degenerate position will produce antibodies tolerant to many, or all, amino acids at the variable
10 positions surrounding one or more fixed residues. Such antibodies will then react with the antigenic determinant in the context of the broadest possible range of surrounding amino acid, peptide, or protein sequences. The fixed
15 residue(s) of the motif may be a single unmodified or modified amino acid, such as a phosphorylated or unphosphorylated residue, or may be multiple unmodified or modified amino acids, such as a consensus recognition site.

As used herein, "antibodies" means polyclonal or
20 monoclonal antibodies, including Fc fragments, Fab fragments, chimeric antibodies, or other antigen-specific antibody fragments.

As used herein, "motif-specific, context-independent
25 antibodies" means antibodies which are specific against one or more fixed amino acid residues in the context of variable surrounding peptide or protein sequences; such antibody

specificity is thus highly independent of the context in which the antigen occurs.

As used herein, "substrate" means any target molecule, including peptides or proteins, which an enzyme specifically recognizes and acts upon.

The general method by which motif-specific, context-independent antibodies are produced in accordance with the present invention is as follows:

(1) Motif-specific antibodies that react with any protein or peptide containing specific target residues independently of the surrounding amino acids may be obtained by synthesizing a highly degenerate peptide library. In one preferred embodiment, the library comprises XXXXXXJ*XXXXXXC where X= all 20 amino acids except cysteine and J*= a modified (*) amino acid (J), for example, phosphothreonine (T*) or acetylated-lysine (K*). It will be appreciated that the specific target residue may be unmodified and that a shorter or longer library may be generated and less than all of the surrounding amino acids may be varied. In one preferred embodiment, the peptide library is about 6 to 14 residues long. While the preferred embodiment utilizes one fixed amino acid (either modified or unmodified) in a varied surrounding context, other preferred embodiments may utilize a motif comprising several fixed amino acids. Likewise, the

surrounding sequence of the library may be varied at more than one position simultaneously, or, as in the preferred embodiment, varied at only one surrounding sequence position per degenerate molecule, such that a library is produced which
5 is completely degenerate at every position except the fixed residue(s). The peptide library can be synthesized by standard F-Moc solid phase peptide synthesis using an ABI peptide synthesizer and using mixtures of each amino acid during degenerate coupling reactions.

10

The incorporation of modified amino acids at fixed positions should not be limited to phosphorylation or acetylations as other modified protected amino acids can also be incorporated, for example, amino acids modified with lipids
15 (e.g. farsynlated, isoprenylated) or protected O-linked or N-linked sugars (e.g. glycosylated), methylated, or ribosylated amino acids, or nucleotides, polymers of nucleotides, nucleosides, or amino acids such as ubiquitin, or amino acid analogues.

20

The incorporation of unmodified amino acids at fixed positions may be selected to mimic conserved motifs, for example zinc fingers or repeating arginine residues.

25

(2) In order to produce as equal a representation of each amino acid as possible at each degenerate position, several rounds of altering the amino acid composition, synthesizing,

and peptide sequencing are conducted. Amino acid sequence analysis at several different positions along the peptide is conducted to verify a random amino acid representation at each position and that the random representation is

5 maintained throughout the synthesis. It will be recognized by one of skill in the art that the number of rounds may vary in order to achieve an equal distribution of all amino acids at each position.

10 (3) The highly diverse peptide library is used as an antigen, preferably by covalent coupling to a carrier. In a preferred embodiment, keyhole limpet hemocyanin (KLH) emulsified in Freund's adjuvant is used as the coupling agent, and the coupled peptide library injected intradermally into a

15 host, such as female New Zealand white rabbits. Booster injections may be given in incomplete Freund's adjuvant until an immune response is obtained. Antibody titre is measured by a suitable method, such as ELISA against the motif-specific peptide libraries. Antisera raised in this manner may

20 be used in both crude or purified preparations, as outlined below.

(4) Antisera from the most promising hosts are purified, for example over protein A, and adsorbed over a J

25 (non-modified) peptide library column. In the preferred embodiment, the nonadsorbed fraction (flow through) is then applied to a J* column, eluted at suitable pH, dialyzed and

tested for J* specificity by a suitable method, such as ELISA using J* and J as antigen.

(5) Antibodies affinity purified in this fashion

5 recognize the J* peptide library but do not react with the J library and exhibit a high degree of specificity for J*. These antibodies may be further tested for lack of reactivity against the unmodified form of the target modified amino acid, J*, or a J* homologue, utilizing a suitable method, such as ELISA.

10

(6) Antibodies may be further tested by western blotting, or another suitable method, using cell extracts prepared from cells treated with and without a selected protein modification enzyme inhibitor, such as protein

15 phosphatase inhibitor okadaic acid. Treatments that increase protein modification will increase the number of antibody reactive proteins as well as the intensity of reactivity. The J* specific antibodies will react with a relatively small number of proteins from control extracts but will react with a
20 very large number following treatment with the selected inhibitor. The antibodies will show no reactivity with the inactive-non-modified versions of these proteins, demonstrating a high degree of J* specificity and suggesting broad cross-reactivity to many different modified -target
25 containing proteins.

(7) The degree of context-independence may be more

carefully examined, for example, by ELISA analysis against individual J* peptides that are mixed together or tested individually. Such analysis can indicate if poor reactivity occurs with certain motifs, such as when J* is followed by
5 proline, for example.

(8) The context-dependence of J* antibody recognition may be further examined, as in the preferred embodiment, using a immobilized grid of modified-peptide libraries. In
10 addition to a fixed target residue, J*, each different library is synthesized to contain an additional fixed amino acid at different positions relative to J* but with all other positions containing all 20 amino acids except cysteine. Each peptide library is coated, for example, on the bottom of an ELISA well
15 and exposed to the J* antibodies. Antibodies that do not react with a particular spot (peptide library) on the grid do not bind when the specified amino acid is present at the specified position. This analysis determines whether or not a particular amino acid at a particular position relative to J* will allow or
20 block binding.

Alternatively, purified antibodies can be linked to beads, allowed to bind the modified or unmodified library, unbound sequences washed away, and bound sequences recovered and
25 subject to amino acid sequencing to determine the amount of each amino acid present at each position in the library. This information will indicate what amino acids are tolerated at

each position.

(9) Monoclonal antibodies may be prepared, as in one form of the preferred embodiment, by coupling the J* peptide library to a suitable carrier, such as KLH, and injected into a host, such as balbC mice. The J* peptide-KLH conjugate may be emulsified in Freund's adjuvant and booster injections in incomplete Freund's adjuvant may be carried out every other week until a response is obtained.

10

(10) Antibody titre is measured by a suitable method, such as ELISA against J* and non-J* peptide libraries. Sera from hosts showing high-titre responses are adsorbed with immobilized non-J* peptide and the nonadsorbed fraction tested by, for example, western blotting.

15

(11) Spleens from hosts showing J*-specific responses are fused to myeloma cells and hybridoma clones are selected and screened. Supernates from individual clones are screened first for their ability to bind the J*-peptide library. Positive clones are next screened for their cross-reactivity against the non-J* library. Clones showing the highest degree of J*-specificity are chosen for further analysis as described above in steps (5) through (8).

20

(12) Overproduction of monoclonal antibodies resulting from step (11) above may be carried out, for example, by

harvesting ascites, culturing selected hybridoma clones, or cloning into a host organism, such as *E. coli*.

The motif-specific, context-independent antibodies
5 produced by this method may be used to identify an unknown
substrate of an enzyme. Such antibodies are first generated
against a motif that is recognized by the enzyme of interest,
for example, a consensus site. These antibodies are then used
to screen a sample for the presence of other, unknown
10 substrates which contain the same motif. This method
enables the rapid detection of important new substrates in a
variety of cascades which involve conserved substrate motifs.
For example, antibodies that selectively recognize a wide
variety of proteins only when phosphorylated at the MAPK
15 consensus phosphorylation site would greatly facilitate the
detection of new MAP kinase targets. MAP kinase could be
overexpressed in cell culture, activated by growth factors,
and target substrate proteins identified by western blotting
using antibodies that selectively recognize the phosphorylated
20 substrate proteins (Stukenberg et al., *Curr. Biol.* 7:338-348
(1997). Alternatively, MAPK could be used to phosphorylate
cDNA expression libraries *in vitro* and MAPK consensus-site
antibodies used to identify cDNA clones expressing MAPK
phosphorylated substrates (Funkunaga and Hunter, *EMBO*
25 16(8):1921-1933 (1997).

Similarly, antibodies produced by the method of the

instant invention may be used to identify an enzyme which modifies a known substrate motif. Such antibodies, whether specific for modified (e.g. phosphorylated) or unmodified (e.g. zinc finger) motifs, can be used to detect whether a certain
5 enzyme of interest has modified a substrate which contains that motif. This method allows for the rapid detection of important new proteins which act on known classes of substrates containing conserved motifs, for the example MAPK consensus site.

10

The motif-specific, context-independent antibodies of the instant invention may also be used *in vitro* as reagents in high-throughput assays, such as drug screens, to detect the enzymatic modification of certain substrates containing a
15 conserved motif. For example, antibodies specific for a certain phosphorylated motif enable the rapid detection of inhibitors of the enzyme that act at that motif. In the case of a drug screen, a single motif-specific antibody can be used to assay the activity of a wide range of enzymes acting at many
20 diverse sequence motifs. Phosphotyrosine antibodies are currently employed in high throughput kinase assays to screen for selective, high affinity tyrosine kinase inhibitors. Compounds or drugs that block enzyme activity are detected by their ability to inhibit kinase activity as determined by a
25 reduction of phosphotyrosine antibody binding to phosphorylated substrate. Similar assays can be set up to screen for pharmaceutically useful compounds using

antibodies produced as described above for phosphoserine, phosphothreonine, or antibodies detecting other protein modifications.

- 5 Antibody based detection of protein kinase activity has several advantages over radioactive assays for use in automated high throughput kinase assays. First, radioactive assays are difficult to automate because they employ transfer of ^{32}P gamma-labeled ATP to a peptide substrate. The
- 10 phosphopeptide is then separated from labeled ATP using phosphocellulose filters and several washing steps, and finally, phosphorylation is quantitated by liquid scintillation methods. Together these steps are time consuming and difficult to automate. Antibody detection allows a wide
- 15 variety of ELISA-type assays that are well suited for automation and high throughput screens.

- Second, radioactive assays require low levels of ATP to insure high levels of ^{32}P incorporation for maximal
- 20 sensitivity. Low levels of ATP in the kinase assay bias the search for inhibitors towards compounds that compete with ATP binding in the protein kinase catalytic cleft. Such screens consistently yield competitive inhibitors at the ATP binding site which due to the highly conserved nature of this
- 25 binding site results in inhibitors with poor selectivity.

Current high-throughput kinase assays typically utilize

biotinylated peptide substrates immobilized on the bottom of a 96 or 386 well plate that is subsequently incubated together with the desired protein kinase, ATP, and the appropriate kinase buffer. Kinase activity is measured using a

5 fluorescently labeled phosphospecific-antibody that reacts only with the phosphorylated peptide substrate. These assays come in two formats homogeneous (not involving wash steps and heterogeneous (involving wash steps). Homogeneous fluorescent assays typically utilize lanthanide-labelled

10 phosphoantibody binding to a phosphorylated peptide substrate that has linked to it an energy acceptor, for example allophycocyanin. Binding of the phosphoantibody to the phosphorylated peptide substrate brings the two fluorophores close enough together to allow fluorescence resonance energy

15 transfer to occur shifting the frequency of the emitted signal, indicating the presence of a biomolecular complex. Different compounds are added to each well and the ability of the compound to inhibit substrate phosphorylation is determined by inhibition of fluorescence energy transfer. This

20 format is similar to the scintillation proximity assay commonly used in radioactive assays. Other homogeneous assays involve the use of fluorescence polarization to measure the binding of phosphoantibody to phosphorylated substrate.

25

The key feature in the homogeneous assays are the limited number of steps and the ease in automation. A large

variety of heterogeneous kinase assays based upon ELIZA formats are also currently in use. These assays typically utilizing fluorescently labeled phosphoantibodies binding phosphorylated peptide substrates that are immobilized in 96
5 or 386 well formats. In this case wash steps are required to separate bound from unbound antibody. Fluorescently labeled antibody retained in the well is then detected using time resolved fluorescence.

10 The motifs used to generate antibodies for such modification screening assays may be either modified or unmodified substrate motifs. Antibodies generated against unmodified motifs will not bind if the substrate has been subsequently modified by an enzyme. Similarly, antibodies
15 generated against modified motifs can detect increases in modified substrate concentrations owing to enzymatic activity.

Similar approaches may be applied to study a variety of
20 other enzymatic modifications, and are not limited to the protein kinase or acetyltransferase activities discussed below. For example, the approach could be used to generate antibodies that recognize many other types of protein modification, including, but not limited to, the addition of
25 sugars, methyl groups, carboxyl groups, the addition of various lipids, or the addition of nucleotides, or polymers of nucleotides, nucleosides, or amino acids such as ubiquitin.

Likewise, such motif-specific, context-independent antibodies may be used on a genome-wide scale to simultaneously profile large and diverse protein populations which contain conserved motifs. A specific two or three amino acid binding site, for example consecutive arginine residues, should appear (based upon a random distribution of amino acids) once every 400 or 8000 residues, respectively, (equating to approximately once per protein, or once every 20 proteins, respectively, (assuming the average protein is 400 amino acids)). Thus, an antibody specific for such a motif independent of the context in which it occurs allows for the rapid screening of a great number of proteins.

Phosphorylation specific antibodies allow genome wide profiling of changes in phosphorylation of proteins as a result of drug treatment or the overexpression of specific genes/proteins as a result of such treatment. Such antibodies also facilitate the profiling of expression of specific proteins in sequenced genomes.

For example, suppose that a drug is developed which inhibits the cell-cycle dependent protein kinase cdc2. The drug has been shown to inhibit cdk2 with high affinity, but the specificity of the compound needs to be further tested to examine whether other protein kinases are inhibited and if so, which ones.

As an early step in this process cell lines may be treated with the drug and the effects on total cell protein phosphorylation monitored using a panel of motif-specific and
5 general phosphoantibodies to examine the nature of the phospho-substrates inhibited by the compound or lead drug.

Total protein from cell extracts prepared from control or drug treated cells may be fractionated using, for example,
10 2-dimensional gels (isoelectric focusing in the first dimension and standard SDS-polyacrylamide molecular weight fractionation in the second dimension), transferred to nitrocellulose membranes, and analyzed by western blotting using, in this hypothetical case, kinase consensus site-
15 specific phosphoantibodies.

In this case, global analysis of total cell proteins using a cdc2 consensus site specific antibody would provide information regarding the ability of the drug to block
20 phosphorylation at all potential cdc2 site substrates. The pattern of inhibition at other non-cdc2 substrates (i.e. the degree of specificity) could also be examined using antibodies to different kinase consensus sites, or using antibodies to phosphotyrosine to determine whether the inhibitor also acts
25 to block tyrosine kinases.

Currently, for mammalian cells, the the identity of the

majority of protein "spots" visualized on 2-D gels are unknown. However, as all human genes are identified and sequenced and the corresponding proteins characterized and "spots" identified, analysis by protein profiling in accordance
5 with the present invention will become even more powerfully informative. The identity of the proteins inhibited will not only confirm the drug specificity but the identity of additional "nonspecific" proteins inhibited will also suggest possible side effects. Identical analysis can be carried out in simpler,
10 completely sequenced organisms, such as yeast where many of the protein "spots" on 2-D gels have already been identified.

The Examples presented below are only intended as specific preferred embodiments of the present invention and
15 are not intended to limit the scope of the invention except as provided in the claims herein. The present invention encompasses modifications and variations of the methods taught herein which would be obvious to one of ordinary skill in the art.

20

The references cited above and below are herein incorporated by reference.

EXAMPLE 1

25 **Context-Independent Phosphothreonine Antibodies:**
Synthesis of peptide library antigens:

Phospho-specific antibodies that react with any protein

containing phosphorylated threonine residues, i.e that bind phosphothreonine independently of the surrounding amino acids, were obtained by synthesizing a highly degenerate peptide library XXXXXXThr*XXXXXXC where X= all 20 amino acids except cysteine and Thr*=phosphothreonine.

The phosphothreonine peptide library was synthesized by standard F-Moc solid phase peptide synthesis using an ABI peptide synthesizer and using mixtures of each amino acid during degenerate coupling reactions. Degenerate peptides were synthesized using an ABI model 433A peptide synthesizer, using FastMoc chemistry (Fields et al., *Pept. Res.* 4:95-101 (1991), hereby incorporated by reference herein) at a scale of 0.085 mmol. Fmoc/NMP chemistry utilizing HBTU amino acid activation (Dourtoglou et al., *Synthesis* 1984: 572-574 (1984), Knorr et al., *Tetra. Let.* 30:1927-1930 (1989), Knorr et al., in Peptides 1988 37-129 (1989), Walter de Grueter & Co, all hereby incorporated by reference herein) was employed for all cycles. Preloaded Fmoc-Cys(Trt) HMP (p-hydroxymethylphenoxyethyl) polystyrene resin functionalized at 0.5 mmol/g was used for each degenerate pool of peptides. Peptides were synthesized using single coupling during each cycle, although coupling times were extended at each position containing a phosphorylated amino acid. The final Fmoc was removed during synthesis. Utilization of preloaded HMP resin along with final Fmoc group removal yields peptides having both free amino and carboxy

termini after cleavage and deprotection.

In order to produce as equal a representation of each amino acid as possible at each degenerate position several rounds of altering the amino acid composition, synthesizing, and peptide sequencing were conducted. The desired peptide pools were to contain an equimolar mix of 19 amino acids (all standard amino acids except Cys) at each "degenerate" site. Because the rate of reactivity of each protected amino acid differs, simply mixing equimolar amounts (each at approximately 5.26% of total) does not result in a population of peptides that is equimolar at each position. In order to maximize degeneracy at each residue, peptide synthesis was first done using equimolar "mixes" at each position.

Phenylthiocarbamyl-amino acid analysis was performed therefore allowing assessment of relative amino acid content at each position. Based on amino acid analysis the molar amounts of each amino acid in the "mix" were adjusted to compensate for different reaction rates, in order to ensure equal representation of each amino acid at each degenerate position. Several rounds of peptide synthesis followed by amino acid analysis were necessary to optimize the amino acid mix, which resulted in a totally degenerate peptide. The optimized amino acid mix arrived at was as follows: G (4.6%); A (5.6%); V (3.3%); L (2.5%); I (4.25%); S (4.4%); T (8.4%); F (2.25%); Y (6.0%); W (6.8%); M (2.9%); P (2.5%); D (5.8%); N (9.5%); E (6.2%); Q (9.4%); K (6.1%); R (6.4%); H (3.5%).

Cleavage of the degenerate peptides from the resin along with removal of side chain protecting groups occurs simultaneously upon treatment with TFA. The cleavage mixture (Perkin Elmer, Emerville, CA (1995)) consists of the following: 0.75g phenol, 0.125 ml methyl sulfide, 0.25 ml 1,2-ethanedithiol, 0.5 ml milliQ H₂O, 0.5 ml thioanisol, 10 ml TFA. The entire mixture was added to the peptide resin (approx. 300 mg). The resin was flushed with nitrogen and gently stirred at room temperature for 3 hours. The resin was then filtered allowing the peptide to be precipitated into cold (0°C) methyl-t-butyl ether. The ether fraction was centrifuged allowing collection of the precipitate. The peptide precipitate was vacuum dried, analyzed by mass spectroscopy, and HPLC purified.

A sample of the peptide was dissolved in acetonitrile/water (50:50, v/v) and analyzed on a Perceptive Biosystems (Framingham, MA) MALDI-TOF mass spectrometer using 2,4,6-trihydroxyacetophenone plus ammonium citrate as the matrix. As expected, the peptide mixture did not show a homogeneous product. MALDI-TOF analysis demonstrated that the peptide pool was degenerate, showing an average mass and the expected statistically normal curve of peptide mass.

25

Peptides were purified using a Waters HPLC system consisting of a Lambda-Max Model 481 Multiwavelength

detector, 500 series pumps, and Automated gradient controller. A Vydac semi-preparative C18 column was used for reverse-phase purification. A 60 min. linear gradient, 10%-100% B, was used at a flow rate of 2 ml/minute. Buffer
5 A consisted of 0.1% TFA/H₂O (v/v) while buffer B consisted of 0.1% TFA/60% CH₃CN/40% H₂O (v/v/v). Detection was at 214 nm.

Because the peptide pool was degenerate (as
10 demonstrated by mass spectroscopy) HPLC purification was not expected to yield a homogeneous product. Base-line separation of peptide mixtures was not achieved by this method and it was only intended as a crude purification/desalting step. Mass spectroscopy was
15 performed and all fractions whose mass was within the theoretical range were pooled and lyophilized.

Amino acid sequence analysis at several different positions along the peptide indicated a random amino acid
20 representation at each position and that the random representation was maintained throughout the synthesis. The results indicated the production of highly diverse peptide libraries that would serve as suitable antigens.

25 Production of Rabbit polyclonal antibodies:

All peptides synthesized contained C-terminal cysteine residues allowing conjugation to the carrier protein (KLH)

using the heterobifunctional cross-linking reagent m-maleimidobenzoyl-N-hydroxysuccinimide ester (MBS). The conjugation procedure used was as described by the manufacturer (Pierce), although the amount of peptide coupled
5 to KLH was increased to 10 mg in order to provide increased material for immunization and boosting of animals. Scale-up required use of a larger desalting column (Bio-Rad 10 DG (Cambridge, MA)) to remove the excess MBS after reaction to N-termini and the ϵ -amino group of KLH Lysine residues.

10

The phosphothreonine peptide library was covalently coupled to keyhole limpet hemocyanin (KLH) (250 μ grams), emulsified in Freund's adjuvant and injected intradermally into female New Zealand white rabbits. Booster injections
15 (200 μ grams) in incomplete Freund's adjuvant were carried out every other week until a response was obtained. Rabbit sera was screened at three week intervals for the presence of phosphopeptide specific immunoreactivity by ELISA using both the phosphothreonine and nonphosphothreonine peptide
20 libraries. When the titre of antibody against phosphopeptide reached 10^5 , rabbits were put on a production bleed schedule with bleeds collected every two weeks. When 40 ml of high titre serum were obtained, purification of phosphospecific antibodies was initiated, as described below.

25

Antisera from the most promising rabbit was purified over protein A and passed over a nonphospho Thr/Ser peptide

library column. The nonadsorbed fraction (flow through) was applied to a phosphothreonine column, eluted at low pH, dialyzed and tested for phosphospecificity by ELISA using phospho- and nonphosphopeptides. Antibodies affinity-
5 purified in this fashion recognized the phosphorylated threonine peptide library but did not react with the nonphosphothreonine/serine library, indicating a high degree of specificity for phosphothreonine (see Figure 1a). ELISA results also indicated that the antibodies also reacted
10 specifically with a mixture of 18 different phosphothreonine peptides but showed no reactivity with any of the corresponding nonphosphopeptides (Figure 1b). The antibodies also exhibited a strict preference for phosphothreonine, showing no reactivity with a mixture of 38 different
15 phosphoserine peptides (Figure 1b) or peptides containing phosphotyrosine.

We next tested the antibodies by western blotting using cell extracts prepared from cells treated with and without the
20 protein phosphatase inhibitor okadaic acid. As shown in Figure 1c the phosphothreonine antibodies react with a relatively small number of proteins from control extracts but react with a very large number following treatment with okadaic acid (see the smear of high Mol Wt. reactive proteins
25 in Figure 1c, lane 2). The antibodies also reacted specifically with the active forms of MAPK (ERK1) and MKK3 only when phosphorylated at threonine residues at their respective

activation loops. The antibodies showed no reactivity with the inactive-nonphosphorylated versions of these proteins (Figure 1c, lanes 3-6). These results demonstrate a high degree of phosphothreonine specificity and suggest broad cross-reactivity to many different threonine-phosphorylated proteins and peptides.

To examine more carefully the degree of context-independence, ELISA analysis was conducted against individual threonine phosphorylated peptides that were mixed together in the previous experiment. As shown in Figure 1a, the phosphothreonine antibody reacts well with all phosphopeptides except those where phosphothreonine is immediately followed by proline, for example the c-Myc and APP1 phosphopeptides (Figure 2b). These results indicate that purified rabbit antibodies reacted in a phosphospecific manner with a wide variety of phosphothreonine but react only poorly with phosphopeptides where the phosphorylated threonine is followed by proline.

20

The context-dependence of phosphothreonine antibody recognition was further examined using a immobilized grid of phosphopeptide libraries. In addition to a fixed phosphothreonine, each different library was synthesized to contain an additional fixed amino acid at the -4, -3, -2, -1, +1, +2, +3 positions relative to phosphothreonine but with all other positions containing all 20 amino acids except cysteine.

Each peptide library was coated on the bottom of an ELISA well and exposed to the phosphothreonine antibodies.

Antibodies that do not react with a particular spot (peptide library) on the grid do not bind when the specified amino acid
5 is present at the specified position. This analysis determines whether or not a particular amino acid at a particular position relative to phosphothreonine will allow or block binding (Figure 1d).

10 Results confirmed that the phosphothreonine antibodies tolerated all amino acids in the -1, -2, -3, -4, and +2, +3 position, and bound equally well to every amino acid except proline at the +1 position (see Figure 1d, first row). The reactivity as defined by this binding profile indicates that the
15 antibodies will bind all phosphothreonine containing sequences except those followed immediately in -1 position by proline. Further analysis using a variety of specific phosphothreonine containing peptides confirmed these results.

20 Phosphothreonine specific antibodies from several other rabbits immunized with the same peptide library antigens were further purified and characterized. Antibodies purified from sera obtained from two other rabbits also produced broadly cross-reacting phosphothreonine antibodies as
25 determined by ELISA. One rabbit produced antibodies that react equally well with peptides containing proline following the phosphothreonine. Taken together, these results

demonstrate the broad context-independence of the phosphothreonine response obtained when combinatorial peptide libraries are used as immunogens.

5

EXAMPLE II

Protein Kinase Consensus Site-Specific Phosphoantibodies:

MAPK-consensus recognition sites: PXS*P

10 A peptide library of the preferred site for MAPK phosphorylation PXS*P was synthesized (Figure 2a) substantially as described in Example I. In addition to an equimolar mix of phosphoserine and threonine, amino acids at two other positions were also fixed; proline at -2 and proline
15 at +1. This library was coupled to KLH and injected into rabbits as described for phosphothreonine. IgG from the most promising rabbit was protein A purified and passed over a nonphospho-Thr/Ser peptide library column. The nonadsorbed fraction (flow through) was applied to a phospho-PXS*P
20 column, eluted at low pH, dialyzed and tested for phosphospecificity by ELISA using phospho- and nonphosphopeptides.

Antibodies affinity purified in this fashion reacted
25 strongly with the phosphorylated PXS*P peptide library but did not react with the nonphosphothreonine/serine library (see Figure 2a). ELISA results also indicated that the antibodies also reacted specifically with a mixture of 18 different

phosphothreonine peptides but showed no reactivity with any of the corresponding nonphosphopeptides (Figure 2a). In addition to being phosphospecific, the antibodies exhibited a preference for proline at the -2 and +1 positions and showed
5 no reactivity with phosphorylated peptides that lack proline at this position (Figure 2a). The antibodies reacted strongly with the RB and cdk4 phosphopeptides but showed no reactivity with the MKK3, PKC α , or p70S6 phosphopeptides that lack proline at the +1 position (Figure 2a). These
10 antibodies do react with some peptides lacking proline at -2, for example the cdk4 phosphopeptide, suggesting that proline at this position is not absolutely necessary.

PXS*P antibodies were further tested by western
15 blotting using cell extracts prepared from cells treated with and without the protein phosphatase inhibitor okadaic acid. Binding of the PXS*P antibodies to cell extracts from RS 4;11 cells was strongly enhanced following treatment with okadaic acid (smear of high Mol Wt. proteins in Figure 2b, lane 2). The
20 antibodies also reacted specifically with ATF-2 phosphorylated in vitro with MAP kinase but not the nonphosphorylated form of this protein (Figure 2b, lanes 3 and 4), demonstrating a high degree of phospho-specificity and broad cross-reactivity to many different phosphorylated
25 proteins and peptides.

The specificity of PXS*P antibody recognition was also

- 37 -

examined using an immobilized grid of phosphopeptide libraries. As described above, in addition to a fixed phosphothreonine or phosphoserine, each different library was synthesized to contain an additional fixed amino acid at the -
5 1, +1, +2 positions relative to phosphothreonine but with all other positions containing all 20 amino acids except cysteine.

The PXS*P antibody reacted weakly with peptide libraries where proline was fixed at the -1 position and
10 reacted strongly with libraries where proline was fixed at both the -2 and +1 positions. The reactivity as defined by this binding profile indicates that the PXS*P antibodies strongly bind only sequences containing the PXS*P motif, as expected, but that the antisera still contain some residual reactivity to
15 S*P (as a result of impurities), which could be removed by further purification using immobilized S*P peptide library.

EXAMPLE III

20 Protein Kinase Consensus Site-Specific Phosphoantibodies:

14-3-3 binding site: RSXS*XP

Antibodies that identify 14-3-3 targets were obtained by synthesizing a peptide library: XXXXRSXS*XPXXXXC where S*
25 is phosphoserine and X represents any amino acid and C is cysteine. The above 14-3-3 phosphopeptide library was synthesized by standard F-Moc solid phase peptide synthesis using an ABI peptide synthesizer and mixtures of each amino

acid except cysteine during degenerate coupling reactions, as discussed in Example I.

The 14-3-3 phosphopeptide library was coupled to KLH
5 and injected into rabbits as described above for
phosphothreonine and PXS*P. Antisera from the most
promising rabbit was purified over protein A and adsorbed
over a nonphospho-14-3-3 peptide library column. The flow-
through of this column was applied to a phospho-14-3-3
10 column eluted at low pH, dialyzed and tested for
phosphospecificity by ELISA using phospho-and nonphospho-
14-3-3 peptide libraries. These affinity purified phospho-14-
3-3 antibodies recognized the phosphorylated 14-3-3 peptide
library but not the nonphospho-14-3-3 library, indicating a
15 high degree of specificity for phospho-14-3-3 (see Figure 3a).
The antibodies also reacted strongly with several different
peptides containing the 14-3-3 motif including; phospho-Bad-
Ser136, cdc25-Ser216, and more weakly with phospho-Bad-
Ser112 which contains a slight variant motif. The antibodies
20 showed no reactivity with the corresponding nonphospho-
peptides (Figure 3a) or with many other phosphopeptides that
did not contain the motif.

Phospho-14-3-3 antibodies were further tested by
25 western blotting using cell extracts prepared from cells
transfected with a GST-Bad fusion protein and treated with
and without the phorbol ester TPA. The antibodies reacted

with a small number of proteins from control extracts (see Figure 3b). Bad was detected in extracts prepared from transfected cells but not control cells. Since the basal level of Bad phosphorylation is high it was difficult to see
5 increased phosphorylation with TPA, although TPA did induce the phosphorylation of several higher molecular weight proteins (arrow in Figure 3b). These results indicate that the phospho-14-3-3 antibodies can detect phosphorylated Bad and other TPA stimulated phospho-proteins.

10

ELISA analysis against the previously described grid of serine/threonine phosphorylated peptide libraries was also conducted. As expected, the phospho-14-3-3 antibodies have an absolute requirement for proline at the +2 position.

15

EXAMPLE IV

Production of mouse monoclonal antibodies: CDK consensus phosphorylation site PXT*PXR:

20

The PXT*/S*PXR sequence represents a consensus phosphorylation site for many of the cell cycle-dependent proteins kinases (cdks). Antibodies that recognize this phosphorylated motif would be useful to identify new cdk
25 substrates important in controlling cell cycle progression. The PXT*/S*PXR peptide library shown in Figure 4a was coupled to KLH and injected into BALB/c mice. The phosphopeptide-KLH conjugate (50 μ grams) emulsified in

- Freund's adjuvant was injected IP. Booster injections (12.5 to 25 μ grams) in incomplete Freund's adjuvant were carried out every three weeks until a response was obtained. Antibody titre was measured by ELISA against the immunized
- 5 phosphopeptide library. Sera from mice showing high-titre responses were adsorbed with immobilized nonphospho Thr/Ser peptide and the nonadsorbed fraction tested by western blotting (data not shown).
- 10 Splenocytes from a mouse showing phosphospecific responses were fused to myeloma X63Ag8.635 cells (Kearney et al., *J. Immunol.* 123:1548-1550 (1979)) and approximately 1,100 hybridoma clones were selected and screened. Supernates from individual clones were screened first for
- 15 their ability to bind the immunized phosphopeptide library and next for their cross-reactivity against the non-phosphopeptide library. Two different clones showing the highest degree of phospho-specificity were chosen for further analysis. The specificity of clones 6B8 and 5A9 were further
- 20 characterized using the phosphopeptide libraries and phosphopeptides shown in Figure 4a. Both clones reacted specifically with phosphothreonine containing libraries and individual peptides but did not significantly react with phosphoserine containing peptides, indicating that
- 25 phosphothreonine selective clones had been identified. Both clones reacted strongly with peptide libraries where proline is fixed in the -2 and +1 positions relative to

phosphothreonine. Reactivity against T*P and PXT*P libraries does not indicate relaxed specificity since one of 400 and one of 20 peptides in the respective libraries will have the appropriate amino acids at the fixed positions. Both clones
5 reacted strongly with a single RB phosphothreonine peptide containing each of the fixed positions present in the immunized library but did not react significantly with the corresponding nonphosphopeptide.

10 Western analysis shows that okadaic acid treatment of cultured cells dramatically increases the reactivity with both clones 6B8 and 5A9 (Figure 4b). Clone 6B8 is also shown to detect cdc2 phosphorylated RB by western blotting (Figure 4b) but does not react with nonphosphorylated RB protein. Clone
15 5A9 was deposited in accordance with the terms and conditions of the Budapest Treaty on September 4, 1998 with the American Type Culture Collection under ATCC Accession No. HB-12563.

20

EXAMPLE V

Acetylated Lysine Specific Antibodies:

Antibodies specifically reactive against acetylated
25 lysine but not reactive against non-acetylated lysine were obtained by synthesizing the following acetylated lysine peptide library: XXXXXXK*XXXXXXC where K* is acetylated and X

represents any amino acid except cysteine and C is cysteine.
The acetylated lysine peptide library was synthesized as
described previously by standard F-Moc solid phase peptide
synthesis using commercially available fully protected
5 acetylated lysine.

The peptide library was coupled to KLH and injected into
rabbits. The K*-peptide-KLH conjugate (250 μ grams) was
used as immunogen as described for the other phosphopeptide
10 libraries. Antisera from the most promising rabbit were
purified over protein A and adsorbed over a non-acetylated
lysine peptide library column. The flow through of this
column was applied to an acetylated lysine column, eluted at
low pH, dialyzed and tested for phosphospecificity by ELISA.

15

Acetylated-lysine antibodies, affinity purified as
described above, recognized the acetylated lysine peptide
library but not the non-acetylated library, indicating a high
degree of specificity for acetylated lysine as measured by
20 ELISA. The antibodies also reacted specifically with as little
as 0.5 ng of acetylated bovine serum albumin (BSA) but
showed no reactivity with up to 10 μ grams of nonacetylated
BSA (see Figure 5a).

25 The antibodies were further examined by western
blotting using cell extracts prepared from cells treated with
and without anisomycin. The antibodies react with a number of

different proteins present in the C6-cell extracts (Figure 5b). In panels b and c, antibodies were preincubated with 1 μ gram of nonacetylated peptide library (Figure 5b) or 1 μ gram of acetylated peptide library (Figure 5c). Preincubation with

5 nonacetylated peptide library had little effect on antibody reactivity with acetylated control protein or bands visualized in the cell extract (Figure 5c, lanes 5-8). However, preincubation of the antibodies with the acetylated lysine peptide library completely blocked antibody binding to control

10 acetylated BSA as well as binding to many proteins present in the cell extract (Figure 5d, lanes 9-12). These results demonstrate a high degree of specificity for acetylated lysine and indicate that the antibodies recognize a broad spectrum of different sized proteins that contain acetylated lysine in a

15 variety of surrounding sequence contexts (compare Figure 5c and d, lanes 1, 2).

WHAT IS CLAIMED IS:

1. A method for producing motif-specific, context-independent antibodies, said method comprising the steps of:
 - (a) constructing a combinatorial peptide library comprising at least one fixed amino acid and surrounding amino acids, wherein at least one surrounding amino acid is variable; and
 - (b) immunizing a host with said peptide library.
2. The method of claim 1, further comprising the steps of isolating the antisera from said host, and purifying said motif-specific, context-independent antibodies from said antisera.
3. The method of claim 1, further comprising the step of utilizing spleen cells from the host of step (b) to generate at least one monoclonal, motif-specific, context-independent antibody.
4. The method of claim 1, wherein said fixed amino acid is a modified amino acid.
5. The method of claim 4, wherein said modified amino acid is selected from the group consisting of a glycosylated amino acid, a phosphorylated amino acid, an acetylated amino acid, a methylated amino acid, a ribosylated amino acid, an isoprenylated amino acid, a lipid-linked amino acid, and an

amino acid analog.

6. The method of claim 1, wherein said fixed amino acid of step (a) is selected from the group consisting of phosphothreonine, phosphoserine, MAPK consensus recognition sites, 14-3-3 consensus recognition sites, CDK consensus recognition sites, and acetylated lysine.

7. The method of claim 1, wherein said fixed amino acid is an unmodified amino acid.

8. The method of claim 1, wherein said peptide library is between about 6 to 14 amino acids long.

9. A motif-specific, context-independent antibody produced by the method of claims 1, 2 or 3.

10. A motif-specific, context-independent antibody which recognizes at least one fixed amino acid in the context of variable surrounding amino acid or peptide sequences.

11. The motif-specific, context-independent antibody of claim 10, wherein said fixed amino acid is selected from the group consisting of phosphothreonine, phosphoserine, MAPK consensus recognition sites, 14-3-3 consensus recognition sites, CDK consensus recognition sites, and acetylated-lysine.

12. A method for identifying an unknown substrate of an

enzyme, said method comprising the steps of:

(a) generating at least one motif-specific, context-independent antibody against a motif recognized by said enzyme; and

(b) screening a target sample with the context-independent antibody for the presence of an unknown substrate containing said motif.

13. The method of claim 12, wherein the motif of step (a) is selected from the group consisting of phosphothreonine, phosphoserine, MAPK consensus recognition sites, 14-3-3 consensus recognition sites, CDK consensus recognition sites, and acetylated-lysine.

14. A method for detecting the modification state of a substrate, said method comprising the steps of:

(a) generating at least one motif-specific, context-independent antibody against a motif selected from the group consisting of an unmodified substrate motif and a modified substrate motif; and

(b) screening a target sample for the presence of a substrate containing said modification.

15. A method for screening a which inhibits or activates enzyme activity on a substrate, said method comprising the steps of:

(a) generating at least one motif-specific, context-independent antibody against a motif selected from the group

consisting of an unmodified substrate motif and a modified substrate motif; and

(b) screening a target sample treated with said drug for the presence of substrate containing said motif.

16. The method of claim 14 or 15, wherein said motif of step (a) is selected from the group consisting of phosphothreonine, phosphoserine, MAPK consensus recognition sites, 14-3-3 consensus recognition sites, CDK consensus recognition sites, and acetylated-lysine.

17. A method for identifying an enzyme which modifies a known substrate motif, said method comprising the steps of:

(a) generating at least one motif-specific, context-independent antibody against said known substrate motif, wherein said motif is selected from the group consisting of an unmodified substrate motif and a modified substrate motif;

(b) reacting an enzyme sample with said known substrate; and

(c) assaying with the antibody of step (a) for the presence of modified substrate of step (b).

18. The method of claim 17, wherein said motif of step (a) is selected from the group consisting of phosphothreonine, phosphoserine, MAPK consensus recognition sites, 14-3-3 consensus recognition sites, CDK consensus recognition sites, and acetylated-lysine.

19. A method for profiling protein levels in a cell on a genome wide scale, said method comprising the steps of:

- (a) generating at least one motif-specific, context-independent antibody against a conserved substrate motif, wherein said motif is selected from the group consisting of an unmodified substrate motif and a modified substrate motif;
- (b) preparing an extract of said cell; and
- (c) utilizing the antibody of step (a) to profile the levels of one or more proteins containing said motif present in the extract of step (b).

20. The method of claim 19, wherein said motif of step (a) is selected from the group consisting of phosphothreonine, phosphoserine, MAPK consensus recognition sites, 14-3-3 consensus recognition sites, CDK consensus recognition sites, and acetylated-lysine.

21. A method for profiling changes in protein levels in a cell on a genome wide scale which result from drug treatment, said method comprising the steps of:

- (a) generating at least one motif-specific, context-independent antibody against a conserved substrate motif, wherein said motif is selected from the group consisting of an unmodified substrate motif and a modified substrate motif;
- (b) preparing an extract of a cell treated with said drug; and
- (c) utilizing said antibody of step (a) to profile changes in the levels of one or more proteins containing said motif

present in said extract of step (b).

22. The method of claim 21, wherein said motif of step (a) is selected from the group consisting of phosphothreonine, phosphoserine, MAPK consensus recognition sites, 14-3-3 consensus recognition sites, CDK consensus recognition sites, and acetylated-lysine.

FIG. 1A

PEPTIDE	SEQUENCE	1.00E+03	5.00E+03	ANTIBODY DILUTIONS	1.00E+04	6.00E+04	1.00E+05	5.00E+05	1.00E+06
Thr	X-X-X-X-Thr-X-X-X-X-X-Cys	1.02	1.32	0.54	0.34	0.07	0.04	0.02	0.02
Ser-Thr	X-X-X-X-X-Ser-Thr-X-X-X-X-X-Cys	0.11	0.05	0.01	0.00	0.01	0.00	0.00	0.00
Threonine* mix	18 phospho-Thr peptide	1.84	1.13	0.40	0.28	0.10	0.07	0.05	0.05
Serine* mix	38 phospho-Ser peptide	0.12	0.04	0.02	0.02	0.02	0.01	0.00	0.00
Akt-Thr308-P	Ile-Lys-Asp-Gly-Ala-Thr-Met-Lys-Thr-Phe-Cys-Gly-Thr-Pro (SEQ ID NO:16)	1.18	0.65	0.24	0.13	0.03	0.01	0.00	0.00
APPI1-Thr688-P	Asp-Ala-Ala-Val-Thr-Pro-Lys-Lys-Arg-His-Leu-Ser-Lys-Cys (SEQ ID NO:17)	0.14	0.03	0.01	0.01	0.01	0.01	0.01	0.00
C3-P	Asp-Thr-Gln-Ile-Lys-Arg-Asn-Thr-Phe-Val-Gly-Thr-Pro-Phe-Cys (SEQ ID NO:18)	1.71	1.13	0.39	0.22	0.03	0.02	0.02	0.02
CAK-Thr187-P	His-Gln-Val-Val-Thr-Arg-Trp-Tyr-Arg-Cys (SEQ ID NO:19)	1.77	1.15	0.41	0.27	0.06	0.03	0.01	0.01
CAMIV-Thr188-P	His-Gln-Val-Leu-Met-Lys-Thr-Val-Cys-Gly (SEQ ID NO:20)	1.78	1.36	0.63	0.40	0.09	0.05	0.01	0.01
CDC2-Thr187-P	Ile-Pro-Ile-Arg-Val-Tyr-Thr-His-Glu-Val-Val-Thr-Leu-Cys (SEQ ID NO:21)	1.02	0.66	0.14	0.08	0.03	0.01	0.01	0.01
CDK2-Thr159-P	Gly-Val-Pro-Val-Arg-Thr-Tyr-Thr-His-Glu-Val-Val-Thr-Leu-Cys (SEQ ID NO:22)	1.88	1.79	0.51	0.44	0.08	0.04	0.02	0.02
p70S6K-Thr388-P	Asn-Gln-Val-Phe-Leu-Gly-Phe-Thr-Tyr-Val-Ala-Pro-Lys-Lys-Cys (SEQ ID NO:14)	1.99	1.44	0.82	0.39	0.08	0.04	0.01	0.01
PKCalpha-P	Lys-Glu-His-Met-Met-Asp-Gly-Val-Thr-Thr-Arg-Thr-Phe-Cys (SEQ ID NO:15)	1.82	1.63	0.84	0.58	0.16	0.08	0.02	0.02
ERK2-P	Asp-His-Thr-Gly-Phe-Leu-Thr-Glu-Tyr-Val-Ala-Thr-Arg-Trp-Cys (SEQ ID NO:23)	1.56	1.18	0.51	0.30	0.07	0.04	0.02	0.02
Myc 8a58/82-P	Glu-Leu-Leu-Pro-Thr-Pro-Pro-Leu-Ser-Pro-Ser-Arg-Arg-Ser-Cys (SEQ ID NO:24)	0.11	0.05	0.03	0.02	0.02	0.02	0.02	0.02
P38-2P	Leu-Ala-Arg-His-Thr-Asp-Glu-Met-Thr-Gly-Tyr-Val-Ala-Thr-Arg-Cys (SEQ ID NO:25)	0.64	0.30	0.08	0.08	0.04	0.04	0.02	0.02
JNK2P	Ser-Phe-Met-Met-Thr-Pro-Tyr-Val-Val-Thr-Arg-Tyr-Tyr-Arg-Cys (SEQ ID NO:26)	1.48	0.44	0.12	0.07	0.03	0.02	0.02	0.02

FIG. 1B

PEPTIDE SEQUENCE	phospho-Thr Reactivity
XXXXXXS*XXXXXX	—
XXXXY*XXXX	—
XXXXXPXS*/T*PXR/XXXX (SEQ ID NO:27)	++
XXXXRSXS*XPXXXX (SEQ ID NO:28)	—
XXXXRSXSXPXXXX (SEQ ID NO:28)	—
XXXXXPXS*/T*PXXXXX	++
XXXXXPXS/TPXXXXX	—
XXXXT*XXXXXX	+++
XXXXXXS/TXXXXXX	—
21 phospho-Thr peptides mixture	+++
38 phospho-Ser peptides mixture	—
30 phospho-Tyr peptides mixture	—
NEB LIBRARY	
X-X-X-X-D/E-X-X-S*-T*-X-X-X-X-X-C (SEQ ID NO:29)	+++
X-X-X-X-X-X-S*/T*-D/E-D/E-D/E-X-X-X (SEQ ID NO:30)	++
X-X-X-X-F-X-X-F-S*/T*-F/Y-X-X-X-X-C (SEQ ID NO:31)	+++
X-X-X-X-R/K-X-S*/T*-X-X-X-X-X-C (NO:31)	+++
X-X-X-R/K-X-X-S*/T*-X-X-X-X-X-C	+++
X-X-X-X-X-X-S*/T*-F/I/M-X-X-X-X-X-C	+++
X-X-X-X-X-X-S*/T*-F/I-X-X-X-X-X-C	+++
X-X-X-X-X-X-S*/T-P-X-X-X-X-X-C	++
X-X-X-X-X-T*-X-X-X-X-X-C	+++
X-X-X-X-X-P-X-S*/T*-P-X-X-X-X-X-C (SEQ ID NO:1)	++
X-X-X-X-X-X-S/T-X-X-X-X-X-C	—
X-X-X-X-X-P-X-S*/T*-P-X-R/K-X-X-X-C (SEQ ID NO:3)	++
ANTIBODY REACTIVITY	
+++ very strong	ELISA O.D. > 2
++ strong	1 - 2
+ weak	0.2 - 1
- very little	< 0.2

3/11

FIG. 1C




FIG. 1D

Fixed Amino Acid	-5-4-3-2-1 X X X X Ser*/Thr* X X X X X Fixed AA position relative to phospho-Ser*/Thr*							+1+2+3+4+5		
	-4	-3	-2	-1	S*/T*	+1	+2	+3		
Ala	+	+	+	+		+	+	+	+	+
Cys	+	+	+	+		+	+	+	+	+
Asp	+	+	+	+		+	+	+	+	+
Glu	+	+	+	+		+	+	+	+	+
Phe	+	+	+	+		+	+	+	+	+
Gly	+	+	+	+		+	+	+	+	+
His	+	+	+	+		+	+	+	+	+
Ile	+	+	+	+		+	+	+	+	+
Lys	+	+	+	+		+	+	+	+	+
Leu	+	+	+	+		+	+	+	+	+
Met	+	+	+	+		+	+	+	+	+
Asn	+	+	+	+		+	+	+	+	+
Pro	+	+	+	+		+	+	+	+	+
Gln	+	+	+	+		+	+	+	+	+
Arg	+	+	+	+		+	+	+	+	+
Ser	+	+	+	+		+	+	+	+	+
Thr	+	+	+	+		+	+	+	+	+
Val	+	+	+	+		+	+	+	+	+
Trp	+	+	+	+		+	+	+	+	+
Tyr	+	+	+	+		+	+	+	+	+

FIG. 2A

PEPTIDE	SEQUENCE	1.00E+03			5.00E+03			1.00E+04			5.00E+04			1.00E+05			5.00E+05			1.00E+06		
		1.00E+03			5.00E+03			1.00E+04			5.00E+04			1.00E+05			5.00E+05			1.00E+06		
PXSP-P	X-X-X-X-Pro-X-Ser ⁺ -Thr ⁺ -Pro-X-X-X-X-Cys (SEQ ID NO: 10)	1.82			1.97			1.74			1.40			0.70			0.35			0.08		
Threonine mix	18 phospho-Thr peptide mix	1.97			1.37			0.67			0.36			0.13			0.07			0.05		
Ser/Thr	X-X-X-X-X-Ser/Thr-X-X-X-X-X-Cys	0.14			0.03			0.01			0.00			0.00			0.00			0.00		
RB Thr373-P	Val-Ile-Pro-Pro-His-Thr ⁺ -Pro-Val-Alg-Thr ⁺ -Val-Met-Asn-Thr-Cys (SEQ ID NO: 11)	2.07			2.17			1.70			1.20			0.48			0.18			0.03		
MKK3-Thr-P	Ser-Val-Ala-Lys-Thr ⁺ -Met-Asp-Ala-Gly-Cys (SEQ ID NO: 12)	0.06			0.04			0.01			0.00			0.00			0.00			0.00		
PKCalpha-P	Lys-Glu-His-Met-Met-Asp-Gly-Val-Thr-Thr-Aug-Thr ⁺ -Phe-Cys (SEQ ID NO: 13)	0.05			0.02			0.01			0.00			0.01			0.00			0.00		
p70 S6K-Thr389	Asn-Gln-Val-Phe-Leu-Gly-Phe-Thr ⁺ -Tyr-Val-Ala-Pro-Lys-Lys-Cys (SEQ ID NO: 14)	0.11			0.05			0.01			0.00			0.01			0.00			0.00		
cdk4-Thr172-P	Arg-Ile-Tyr-Ser-Tyr-Gln-Met-Ala-Leu-Thr ⁺ -Pro-Val-Val-Val-Lys-Cys (SEQ ID NO: 15)	2.07			2.21			2.01			1.55			0.68			0.31			0.07		

FIG. 2B





FIG. 3A

PEPTIDE	SEQUENCE	ANTIBODY DILUTIONS					1.00E+05	5.00E+05
		1.00E+03	5.00E+03	1.00E+04	5.00E+04	1.00E+05		
14-3-3 BM-P	X-X-X-X-Arg-Ser-X-Ser ⁻ -X-Pro-X-X-X-Cys (SEQ ID NO:6)	2.41	2.15	1.49	1.15	0.44		0.25
14-3-3 BM	X-X-X-X-Arg-Ser-X-Ser ⁻ -X-Pro-X-X-X-Cys (SEQ ID NO:6)	0.07	0.03	0.02	0.03	0.02		0.03
CDC25-Ser216-P	Gly-Leu-Tyr-Arg-Ser-Pro-Ser ⁻ -Met-Pro-Glu-Asn-Leu-Asn-Arg-Cys (SEQ ID NO:7)	2.35	2.08	1.49	1.05	0.33		0.18
CDC25-Ser216	Gly-Leu-Tyr-Arg-Ser-Pro-Ser ⁻ -Met-Pro-Glu-Asn-Leu-Asn-Arg-Cys (SEQ ID NO:7)	0.05	0.02	0.03	0.03	0.04		0.03
Bad-Ser112-P	Thr-Arg-Ser-Arg-His-Ser-Ser ⁻ -Tyr-Pro-Ala-Gly-Thr-Glu-Glu-Cys (SEQ ID NO:8)	1.59	0.43	0.10	0.03	0.01		0.00
Bad-Ser112	Thr-Arg-Ser-Arg-His-Ser-Ser ⁻ -Tyr-Pro-Ala-Gly-Thr-Glu-Glu-Cys (SEQ ID NO:8)	0.00	0.00	0.00	0.00	0.00		0.00
Bad-Ser136	Phe-Arg-Gly-Arg-Ser-Arg-Ser ⁻ -Ala-Pro-Pro-Asn-Leu-Trp-Ala-Cys (SEQ ID NO:9)	0.03	0.00	0.00	0.00	0.00		0.00
Bad-Ser136-P	Phe-Arg-Gly-Arg-Ser-Arg-Ser ⁻ -Ala-Pro-Pro-Asn-Leu-Trp-Ala-Cys (SEQ ID NO:9)	3.25	1.86	0.73	0.51	0.07		0.03

FIG. 3B




9/11

FIG. 4A




PEPTIDE	SEQUENCE	MONOCLONAL ANTIBODIES	
		6B8	5A9
Ser/ThrPro-P	X-X-X-X-X-Ser/Thr-Pro-X-X-X-X-X-Cys	1.774	0.731
ProXSer/ThrPro-P	X-X-X-X-X-Pro-X-Ser/Thr-Pro-X-X-X-X-X-Cys (SEQ ID NO:1)	0.924	0.768
ProXSer/ThrPro-P	X-X-X-X-X-Pro-X-Ser/Thr-Pro-X-X-X-X-X-Cys (SEQ ID NO:2)	0.02	0.063
ProXSer/ThrProXArg-P	X-X-X-X-X-Pro-X-Ser/Thr-Pro-X-Arg/Lys-X-X-X-Cys (SEQ ID NO:3)	1.955	1.275
Thr-P	X-X-X-X-X-X-Thr-X-X-X-X-X-X-Cys	0	--
Ser-P	X-X-X-X-X-X-Ser-X-X-X-X-X-X-Cys	0.031	0.088
Ser/Thr	X-X-X-X-X-X-Ser/Thr-X-X-X-X-X-X-Cys	0.021	0.068
Tyr-P	X-X-X-X-X-X-Tyr-X-X-X-X-X-X-Cys	0.023	0.072
Rb (Ser795)-P	Ser-Pro-Tyr-Lys-Phe-Pro-Ser-Ser-Pro-Leu-Arg-Ile-Pro-Gly-Cys (SEQ ID NO:4)	0.032	0.124
Rb (Thr373)-P	Val-Ile-Pro-Pro-His-Thr-Pro-Val-Arg-Thr-Val-Met-Asn-Thr-Cys (SEQ ID NO:5)	3.336	3.503
Rb (Thr373)	Val-Ile-Pro-Pro-His-Thr-Pro-Val-Arg-Thr-Val-Met-Asn-Thr-Cys (SEQ ID NO:5)	0.02	0.073

10/11

FIG. 4B



11/11

FIG. 5A**FIG. 5B****FIG. 5C****FIG. 5D**

SEQUENCE LISTING

<110> COMB, MICHAEL J.
TAN, YI

<120> Production Of Motif-Specific And Context-Independent
Antibodies Using Peptide Libraries As Antigens

<130> NEB-138

<140>

<141>

<160> 31

<170> PatentIn Ver. 2.0

<210> 1

<211> 15

<212> PRT

<213> Homo sapiens

<220>

<221> PEPTIDE

<222> (8)

<223> X=Ser*/Thr*

<400> 1

Xaa	Xaa	Xaa	Xaa	Xaa	Pro	Xaa	Xaa	Pro	Xaa	Xaa	Xaa	Xaa	Xaa	Cys
1					5				10					15

<210> 2

<211> 15

<212> PRT

<213> Homo sapiens

<220>

<221> PEPTIDE

<222> (8)

<223> X-Ser*/Thr*

<400> 2

Xaa	Xaa	Xaa	Xaa	Xaa	Pro	Xaa	Xaa	Pro	Xaa	Xaa	Xaa	Xaa	Xaa	Cys
1					5				10					15

<210> 3

<211> 15

<212> PRT

<213> Homo sapiens

<220>

<221> PEPTIDE

<222> (8)

<223> X=Ser*/Thr*

<220>

<221> PEPTIDE

<222> (11)

<223> X=Arg/Lys

<400> 3

Xaa	Xaa	Xaa	Xaa	Xaa	Pro	Xaa	Xaa	Pro	Xaa	Xaa	Xaa	Xaa	Xaa	Cys
1					5				10					15

<210> 4

<211> 15

<212> PRT

<213> Homo sapiens

<400> 4

Ser	Pro	Tyr	Lys	Phe	Pro	Ser	Ser	Pro	Leu	Arg	Ile	Pro	Gly	Cys
1				5					10					15

<210> 5

<211> 15

<212> PRT

<213> Unknown

<220>

<220>

<223> Description of Unknown Organism: Unknown

<400> 5

Val	Ile	Pro	Pro	His	Thr	Pro	Val	Arg	Thr	Val	Met	Asn	Thr	Cys
1				5					10					15

<210> 6

<211> 15

<212> PRT

<213> Unknown

<220>

<223> Description of Unknown Organism: Unknown

<400> 6

Xaa	Xaa	Xaa	Xaa	Arg	Ser	Xaa	Ser	Xaa	Pro	Xaa	Xaa	Xaa	Xaa	Cys
1				5					10					15

<210> 7

<211> 15

<212> PRT

<213> Unknown

<220>

<223> Description of Unknown Organism: Unknown

<400> 7

Gly Leu Tyr Arg Ser Pro Ser Met Pro Glu Asn Leu Asn Arg Cys
1 5 10 15

<210> 8

<211> 15

<212> PRT

<213> Unknown

<220>

<223> Description of Unknown Organism: Unknown

<400> 8

Thr Arg Ser Arg His Ser Ser Tyr Pro Ala Gly Thr Glu Glu Cys
1 5 10 15

<210> 9

<211> 15

<212> PRT

<213> Unknown

<220>

<223> Description of Unknown Organism: Unknown

<400> 9

Phe Arg Gly Arg Ser Arg Ser Ala Pro Pro Asn Leu Trp Ala Cys
1 5 10 15

<210> 10

<211> 16

<212> PRT

<213> Unknown

<220>

<223> Description of Unknown Organism: Unknown

<400> 10

Xaa Xaa Xaa Xaa Xaa Pro Xaa Ser Thr Pro Xaa Xaa Xaa Xaa Xaa Cys
1 5 10 15

<210> 11

<211> 15

<212> PRT

<213> Unknown

<220>

<223> Description of Unknown Organism: Unknown

<400> 11

Val Ile Pro Pro His Thr Pro Val Arg Thr Val Met Asn Thr Cys
1 5 10 15

<210> 12
<211> 10
<212> PRT
<213> Unknown

<220>
<223> Description of Unknown Organism: Unknown

<400> 12
Ser Val Ala Lys Thr Met Asp Ala Gly Cys
1 5 10

<210> 13
<211> 14
<212> PRT
<213> Unknown

<220>
<223> Description of Unknown Organism: Unknown

<400> 13
Lys Glu His Met Met Asp Gly Val Thr Thr Arg Thr Phe Cys
1 5 10

<210> 14
<211> 15
<212> PRT
<213> Unknown

<220>
<223> Description of Unknown Organism: Unknown

<400> 14
Asn Gln Val Phe Leu Gly Phe Thr Tyr Val Ala Pro Lys Lys Cys
1 5 10 15

<210> 15
<211> 16
<212> PRT
<213> Unknown

<220>
<223> Description of Unknown Organism: Unknown

<400> 15
Arg Ile Tyr Ser Tyr Gln Met Ala Leu Thr Pro Val Val Val Lys Cys
1 5 10 15

<210> 16
<211> 14

<212> PRT
<213> Unknown

<220>
<223> Description of Unknown Organism: Unknown

<400> 16
Ile Lys Asp Gly Ala Thr Met Lys Thr Phe Cys Gly Arg Pro
1 5 10

<210> 17
<211> 14
<212> PRT
<213> Unknown

<220>
<223> Description of Unknown Organism: Unknown

<400> 17
Asp Ala Ala Val Thr Pro Lys Lys Arg His Leu Ser Lys Cys
1 5 10

<210> 18
<211> 15
<212> PRT
<213> Unknown

<220>
<223> Description of Unknown Organism: Unknown

<400> 18
Asp Thr Gln Ile Lys Arg Asn Thr Phe Val Gly Thr Pro Phe Cys
1 5 10 15

<210> 19
<211> 10
<212> PRT
<213> Unknown

<220>
<223> Description of Unknown Organism: Unknown

<400> 19
His Gln Val Val Thr Arg Trp Tyr Arg Cys
1 5 10

<210> 20
<211> 10
<212> PRT
<213> Unknown

<220>

<223> Description of Unknown Organism: Unknown

<400> 20

His Gln Val Leu Met Lys Thr Val Cys Gly
1 5 10

<210> 21

<211> 14

<212> PRT

<213> Unknown

<220>

<223> Description of Unknown Organism: Unknown

<400> 21

Ile Pro Ile Arg Val Tyr Thr His Glu Val Val Thr Leu Cys
1 5 10

<210> 22

<211> 15

<212> PRT

<213> Unknown

<220>

<223> Description of Unknown Organism: Unknown

<400> 22

Gly Val Pro Val Arg Thr Tyr Thr His Glu Val Val Thr Leu Cys
1 5 10 15

<210> 23

<211> 15

<212> PRT

<213> Unknown

<220>

<223> Description of Unknown Organism: Unknown

<400> 23

Asp His Thr Gly Phe Leu Thr Glu Tyr Val Ala Thr Arg Trp Cys
1 5 10 15

<210> 24

<211> 15

<212> PRT

<213> Unknown

<220>

<223> Description of Unknown Organism: Unknown

<400> 24

Glu Leu Leu Pro Thr Pro Pro Leu Ser Pro Ser Arg Arg Ser Cys

1 5 10 15

<210> 25
 <211> 17
 <212> PRT
 <213> Unknown

<220>
 <223> Description of Unknown Organism: Unknown

<400> 25
 Leu Ala Arg His Thr Asp Asp Glu Met Thr Gly Tyr Val Ala Thr Arg
 1 5 10 15

Cys

<210> 26
 <211> 15
 <212> PRT
 <213> Unknown

<220>
 <223> Description of Unknown Organism: Unknown

<400> 26
 Ser Phe Met Met Thr Pro Tyr Val Val Thr Arg Tyr Tyr Arg Cys
 1 5 10 15

<210> 27
 <211> 14
 <212> PRT
 <213> Unknown

<220>
 <221> PEPTIDE
 <222> (8)
 <223> X=Ser*/Thr*

<220>
 <221> PEPTIDE
 <222> (11)
 <223> X=Arg/Lys

<220>
 <223> Description of Unknown Organism: Unknown

<400> 27
 Xaa Xaa Xaa Xaa Xaa Pro Xaa Xaa Pro Xaa Xaa Xaa Xaa Xaa
 1 5 10

<210> 28

<211> 14
<212> PRT
<213> Unknown

<220>
<223> Description of Unknown Organism: Unknown

<400> 28
Xaa Xaa Xaa Xaa Arg Ser Xaa Ser Xaa Pro Xaa Xaa Xaa Xaa
1 5 10

<210> 29
<211> 15
<212> PRT
<213> Unknown

<220>
<221> PEPTIDE
<222> (5)
<223> X=Asp/Glu

<220>
<223> Description of Unknown Organism: Unknown

<400> 29
Xaa Xaa Xaa Xaa Xaa Xaa Xaa Ser Thr Xaa Xaa Xaa Xaa Xaa Cys
1 5 10 15

<210> 30
<211> 13
<212> PRT
<213> Unknown

<220>
<221> PEPTIDE
<222> (7)
<223> X=Ser*/Thr*

<220>
<221> PEPTIDE
<222> (8)..(10)
<223> X=Asp/Glu

<220>
<223> Description of Unknown Organism: Unknown

<400> 30
Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
1 5 10

<210> 31
<211> 15
<212> PRT

<213> Unknown

<220>

<221> PEPTIDE

<222> (9)

<223> X=Ser*/Thr*

<220>

<221> PEPTIDE

<222> (10)

<223> X=Phe/Tyr

<220>

<223> Description of Unknown Organism: Unknown

<400> 31

Xaa	Xaa	Xaa	Xaa	Phe	Xaa	Xaa	Phe	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Cys
1				5				10						15

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US99/19597

A. CLASSIFICATION OF SUBJECT MATTER IPC(7) : G01N 33/53; C07K 16/00 US CL : 435/7.1, 4; 530/388.1 According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) U.S. : 435/7.1, 4; 530/388.1 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) CAS ONLINE		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	SONGYANG, Z. et al. A structural basis for substrate specificities of protein Ser/Thr kinases: primary sequence preference of casein kinases I and II, NIMA, phosphorylase kinase, calmodulin-dependent kinase II, CDK5 and Erk1. Molecular and Cellular Biology. November 1996, Vol. 16, No. 11, pages 6486-6493, especially page 6488.	1-8
Y	WESTENDORF, J. et al. Cloning of cDNAs for M-phase phosphoproteins recognized by the MPM2 monoclonal antibody and determination of the phosphorylated epitope. Proc. Natl. Acad. Sci. USA. January 1994, Vol. 91, pages 714-718, especially pages 716-717.	1-8
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents: *A* document defining the general state of the art which is not considered to be of particular relevance *B* earlier document published on or after the international filing date *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) *O* document referring to an oral disclosure, use, exhibition or other means *P* document published prior to the international filing date but later than the priority date claimed	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art *A* document member of the same patent family	
Date of the actual completion of the international search 19 NOVEMBER 1999		Date of mailing of the international search report 03 FEB 2000
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230		Authorized officer T. WESSENDORF Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/19597

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	FAZIOLI, F. et al. Eps 8, a substrate for the epidermal growth factor receptor kinase, enhances EGF-dependent mitogenic signals. The EMBO Journal. 1993, Vol. 12, No. 10, pages 3799-3808, entire document.	1-8
Y	CZERNIK, A. et al. Production of Phosphorylation state-specific antibodies. Methods in Enzymology. 1991, Vol. 201, pages 265-283, entire document.	1-8
Y	HEFFETZ, D. et al. Generation and use of antibodies to phosphothreonine. Methods in Enzymology. 1991, Vol. 201, pages 44-53, entire document.	1-8
Y	FUKUNAGA, R. et al. MNK1, a new MAP kinase-activated protein kinase, isolated by a novel expression screening method for identifying protein kinase substrates. The EMBO Journal. 1997, Vol. 16, No. 8, pages 1921-1922, entire document.	1-8

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/19597

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

Please See Extra Sheet.

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:
1-8, glycosylated species

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US99/19597

BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

This application contains the following inventions or groups of inventions which are not so linked as to form a single inventive concept under PCT Rule 13.1. In order for all inventions to be searched, the appropriate additional search fees must be paid.

Group I, claim(s) 1-8, drawn to a method of producing an antibody.

Group II, claim(s) 9-11, drawn to an antibody.

Group III, claim(s) 12-13, drawn to a method to identify an unknown substrate of an enzyme.

Group IV, claim(s) 14 and 16, drawn to a method to detect a modified substrate.

Group V, claim(s) 15-16, drawn to a method of screening which inhibits or activates enzyme activity.

Group VI, claim(s) 17-18, drawn to a method to identify an enzyme that modifies a known substrate.

Group VII, claims 19-20, drawn to a method to profile a protein.

Group VIII, claims 21-22, drawn to a method to profile a modified protein.

The inventions listed as Groups I and II do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons: the method of Group I can produce a different motif specific antibody as evidenced by .

The inventions listed as Groups I and III-VIII do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons: each of these methods contain recite separate methods that results in different products and/or results and effects.

The inventions listed as Groups II and III-VIII do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons: as the different methods are employed to produce different results.

The inventions listed as Groups III-VIII do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons: as the different methods each contains or employs different process steps that results in different product effects and/or results.

This application contains claims directed to more than one species of the generic invention. These species are deemed to lack Unity of Invention because they are not so linked as to form a single inventive concept under PCT Rule 13.1. In order for more than one species to be searched, the appropriate additional search fees must be paid. The species are as follows:

- 1). Modified amino acids:
 - A). Glycosylated amino acid
 - B). Phosphorylated amino acid
 - C). Acetylated amino acid
 - D). Methylated amino acid
 - E). Ribosylated amino acid
 - F). Isoprenylated amino acid
 - G). Lipid-linked amino acid
 - H). Amino acid analog
- 2). Fixed or Motif amino acids:
 - A). Phosphothreonine
 - B). Phosphoserine
 - C). MAPK consensus
 - D). 14-3-3 consensus
 - E). CDk consensus
 - F). Acetylated lysine

The species lack the same or corresponding special technical features as evidenced by the teachings of e.g., Lewis et al.